



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Stratego® Fungicide

For Control of Wheat Rust, Powdery Mildew,
Leaf Blight, Tan Spot and Glume Blotch in
Wheat.

EPA Reg. No. 264-779

EPA SLN. No. WA-030015

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

This label for Stratego® Fungicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2008.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Wheat

Stratego® will provide effective control of wheat rusts (*Puccinia graminus*, *P. recondita*, *P. striiformis*), powdery mildew (*Erysiphe graminis*), leaf blight (*Septoria tritici*), tan spot (*Pyrenophora tritici-repentis*) and glume blotch (*Stagnospora nodorum*).

Applications may be either by ground equipment or aerially. Stratego® may be applied at any growth stage through Feekes Growth Stage 10.5 (full head emergence). Do not apply after this growth stage to avoid possible illegal residues.

Disease	Rate	Comments
Rusts Powdery Mildew Leaf Blight Tan Spot Glume Blotch	10 fl. oz./A	Begin applications preventively when conditions are favorable for disease development.

RESTRICTIONS: To avoid possible illegal residues:

1. **Do not** exceed more than 10 fluid ounces of Stratego® per acre per season.
2. **Do not** apply within 35 days of harvest.
3. **Do not** graze or feed livestock treated wheat, straw, forage or hay.
4. **Do not** cut the green crop for hay or silage.
5. After harvest, the straw may be used for bedding.
6. Regardless of formulation or product, do not apply more than a total of 1.8 oz (0.1125 lb) active ingredient of propiconazole per acre per season.
7. For use under this SLN label, do not apply this product through any type of irrigation system.

This pesticide is toxic to fish and aquatic invertebrates. Stratego® Fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com